

RESEARCH ARTICLE

# Peripapillary retinal splitting visualized on OCT in glaucoma and glaucoma suspect patients


Dilraj S. Grewal<sup>1a</sup>, Daniel J. Merlau<sup>2b</sup>, Pushpanjali Giri, Marion R. Munk, Amani A. Fawzi, Lee M. Jampol, Angelo P. Tanna\*

Department of Ophthalmology, Northwestern University Feinberg School of Medicine, Chicago, Illinois, United States of America

<sup>1a</sup> Current address: Duke Eye Center, Department of Ophthalmology, Duke University Medical Center, Durham, North Carolina, United States of America

<sup>2b</sup> Current address: Columbia University, New York, New York, United States of America

\* [atanna@northwestern.edu](mailto:atanna@northwestern.edu)



## Abstract

### Purpose

To identify the risk factors for development of peripapillary retinal splitting (schisis) in patients with glaucoma or suspicion of glaucoma

### Setting

Glaucoma Clinic, Department of Ophthalmology, Northwestern University Feinberg School of Medicine, Chicago, IL

### Methods

In this institutional cross-sectional study, 495 patients (990 eyes) who had undergone spectral-domain optical coherence tomography (OCT Spectralis HRA-OCT, Heidelberg Engineering) optic nerve head (ONH) imaging and did not have identifiable optic nerve pits, pseudopits or coloboma were included. OCT scans were reviewed by two observers.

### Main outcome measures

Presence of peripapillary retinal splitting identified on OCT raster scans.

### Results

Eleven of 990 glaucoma and glaucoma suspect eyes (1.1%) of 7 patients (2 females, 5 males, mean age  $64.5 \pm 9.2$  years) had peripapillary retinal splitting. Two of these 11 eyes had extension of the splitting into the macula but none to the fovea. Of these 11 patients, 2 (28.6%) were glaucoma suspects, 3 (42.9%) had primary open-angle glaucoma, 1 (14.3%) had chronic angle-closure glaucoma and 1 (14.3%) had pigmentary glaucoma. 7/11 (63.6%) eyes had vitreous traction to the disc visualized on OCT and 6/11 eyes (54.5%) had beta-zone peripapillary atrophy.

## OPEN ACCESS

**Citation:** Grewal DS, Merlau DJ, Giri P, Munk MR, Fawzi AA, Jampol LM, et al. (2017) Peripapillary retinal splitting visualized on OCT in glaucoma and glaucoma suspect patients. PLoS ONE 12(8): e0182816. <https://doi.org/10.1371/journal.pone.0182816>

**Editor:** Demetrios G. Vavvas, Massachusetts Eye & Ear Infirmary, Harvard Medical School, UNITED STATES

**Received:** March 1, 2017

**Accepted:** July 25, 2017

**Published:** August 23, 2017

**Copyright:** © 2017 Grewal et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** Supported by an unrestricted grant from Research to Prevent Blindness, NY to the Department of Ophthalmology, Northwestern University Feinberg School of Medicine, Heed Ophthalmic Foundation (San Francisco, CA) and Ronald G. Michels Fellowship Foundation (Riderwood, MD) (DSG). None of the authors have

any financial or proprietary interest in this manuscript. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Conclusions

We observed peripapillary retinal splitting in 1.1% of a series of 990 glaucoma and glaucoma-suspect eyes. Evidence of adherent vitreous with traction and peripapillary atrophy was found in a majority of the involved eyes. A comparison to an age and axial length matched cohort is required to determine if this is a condition that is associated with glaucoma.

## Introduction

Peripapillary retinal splitting (retinoschisis), which is characterized by the splitting of the peripapillary retinal nerve fiber layer (RNFL), inner or outer plexiform and/or nuclear retinal layers with schisis cavities, occurs in eyes with congenital cavitory optic disc anomalies (CODA) including, colobomas or pits and morning glory syndrome. [1–4] It has been previously reported in patients with glaucoma without optic disc pits. [5–8] There has also been increased interest in peripapillary and macular retinoschisis in glaucomatous eyes recently. [9–11] The impact of retinoschisis on RNFL thickness measurements has been noted illustrating its effect as a confounding factor in detecting structural progression in glaucoma. [11]

There are no previous reports of the prevalence of peripapillary retinoschisis in glaucoma and glaucoma suspects. In addition, most of the previous reports were based on isolated cases or case series with no identification of factors associated with the presence of peripapillary splitting. In the current study, we used high density optical coherence tomography (OCT) raster scans of the optic nerve head to identify the prevalence of peripapillary splitting, peripapillary atrophy and vitreous traction in a large series of patients with glaucoma or suspicion of glaucoma.

## Methods

The study was approved by the Institutional Review Board at Northwestern University, Feinberg School of Medicine and adhered to the tenets set forth by the Declaration of Helsinki. The Institutional Review Board waived the requirement for written informed consent.

We analyzed the medical records of consecutive patients examined in our glaucoma clinic over a period of 3 months at the Department of Ophthalmology, Northwestern University Feinberg School of Medicine. Exclusion criteria were the presence of CODA, optic nerve pits, pseudopits or coloboma, concurrent retinal disease (for example, vascular disorders like central retinal vein occlusion (CRVO), branch retinal vein occlusion (BRVO), or macular degeneration), optic nerve disease other than glaucoma or media opacity precluding a good quality OCT scan.

Optic nerve head scans of subjects with glaucoma or suspicion of glaucoma underwent spectral domain OCT imaging. The raster lines protocol (4 mm by 4 mm area centered on the optic disc of the Spectralis HRA-OCT was used (Version 4; Heidelberg Engineering, Heidelberg, Germany). The standard raster volume scan (61 B-scans with 125  $\mu$ m inter-scan spacing) was performed. OCT image quality was identified from the quality bar (range 0–40 decibels). Scans with artifacts or scan quality <15 dB were excluded. [12]

OCT data were exported from the imaging instruments and B-scans for each slice were reviewed. Scans with poor quality were excluded. OCT scans for both groups were reviewed by two independent observers. The two observers independently reviewed the individual B scans

for both groups to detect the presence of schisis. The main outcome measure was the presence of peripapillary schisis identified on OCT raster scans. The splitting included any eyes with splitting delineated by cystoid spaces in the retinal nerve fiber layer (RNFL), inner nuclear layer (INL), outer nuclear layer (ONL) and outer plexiform layer (OPL). All cases of schisis were verified by a team of two retina specialists (AAF and LMJ) and one glaucoma specialist (APT). Cases with uncertain findings on the initial evaluation were adjudicated by the same team.

As part of their routine follow-up, each subject underwent a full ophthalmic examination, including assessment of visual acuity, intraocular pressure (IOP) with a Goldmann applanation tonometer, optic nerve head (ONH) evaluation and stereoscopic slit lamp biomicroscopic fundus examination with a 78 or 90-diopter lens, and the 24-2 Swedish Interactive Threshold Algorithm standard automated visual field (VF) test (Humphrey VF Analyzer; Carl Zeiss Meditec, Dublin, California, USA).

## Statistical analysis

Statistical analysis was performed with Statistical Package for the Social Sciences V.20.0 (SPSS, Chicago, Illinois, USA). Data were analyzed descriptively and are given in mean  $\pm$  standard deviation (SD) and in frequencies. Chi square test was used for comparisons between cases of peripapillary splitting among glaucoma and glaucoma suspects and controls. Independent samples t-test was used for continuous variables. *P* values  $<0.05$  were considered statistically significant.


## Results

A total of 990 eyes of 495 consecutive glaucoma patients and glaucoma suspects who had undergone ONH raster scan OCT imaging were included. Peripapillary schisis was present in 11 eyes (1.1%) of 7 patients (2 females, 5 males, mean age  $64.5 \pm 9.2$  years, range 52 to 78 years). On initial review, an additional 4 cases had been identified to have uncertain findings but were determined to be normal by expert consensus. Figs 1, 2 and 3 provide representative examples of eyes identified with peripapillary schisis.

## Clinical characteristics

The clinical and demographic characteristics of all subjects in the study, as well as the associated imaging findings, are summarized in the Table 1. Of the 7 patients with peripapillary schisis, 3 (42.9%) had primary open angle glaucoma (POAG), 2 (28.6%) were glaucoma suspects, 1 (14.3%) had primary angle closure-glaucoma and 1 (14.3%) had pigmentary glaucoma. The distribution of clinical diagnosis among eyes with and without peripapillary schisis (979 eyes of 488 patients) is provided in Table 2. Detailed information on all subjects is provided in S1 Table.

Mean IOP (mean  $\pm$  standard deviation) at the time of imaging for these patients was  $14.4 \pm 3.3$  mmHg as compared to  $16.3 \pm 4.9$  for eyes without peripapillary schisis ( $p = 0.19$ ). For eyes with peripapillary schisis, the mean Humphrey visual field mean deviation (mean  $\pm$  standard deviation) was  $-7.16 \pm 3.32$  dB (median -2.40, interquartile range -0.84 to -7.89) and the mean pattern standard deviation was  $4.69 \pm 4.37$  dB (median 2.16, interquartile range 1.94 to 4.42) which was not significantly different from eyes without peripapillary schisis, at  $-4.93 \pm 7.14$  dB (median 2.13, interquartile range -0.25 to -7.17,  $p = 0.64$ ) and  $4.46 \pm 3.83$  dB (median 2.37, interquartile range 1.73 to 6.43,  $p = 0.51$ ) respectively.



**Fig 1. 78 year old male with primary open angle glaucoma (POAG) with peripapillary retinoschisis.** (1A) Optic nerve photograph shows glaucomatous optic neuropathy (1B) There is mild visual field damage with a mean deviation of -2.19 dB. (1C, 1D and 1E) Three horizontal OCT raster scans through three different sections of the optic nerve demonstrating peripapillary retinal splitting (retinoschisis) with adherent vitreous in the region of retinoschisis (yellow arrow), splitting in the nerve fiber layer (yellow star), and inner plexiform



layer (white star) and outer plexiform layer with a likely outer nuclear layer component as well (white arrows). (1F) Circumpapillary retinal nerve fiber layer (RNFL) thickness map shows retinoschisis nasally within the RNFL segmentation (yellow star), causing a small area of artifactual thickening on the RNFL thickness profile.


<https://doi.org/10.1371/journal.pone.0182816.g001>

## OCT and imaging characteristics

Of the 11 eyes, all 11 (100%) had splitting in the ONL, 8 (72.3%) had splitting in the INL, 7 (63.6%) had splitting in RNFL and 3 (27.3%) had splitting in OPL. Two of these 11 eyes (18.1%) also had extension of the schisis cavity into the macula as visualized on OCT. Seven of the 11 (63.6%) eyes had vitreous traction in the peripapillary area visualized on OCT and 6/11 eyes (54.5%) had beta-zone peripapillary atrophy. In contrast, among the 979 eyes without retinoschisis, 13 (1.33%) had vitreous traction in the peripapillary area visualized on OCT ( $p < 0.001$ ) and 245 eyes (25.03%) had beta zone peripapillary atrophy ( $p = 0.18$ , Chi-Square test). Eight of the 11 eyes with schisis (72.7%) had glaucomatous optic disc excavation.


## Axial length and refractive status

Among eyes with available axial length measurements ( $n = 85$  eyes), mean axial length (mm) was  $25.22 \pm 2.54$  among eyes with peripapillary schisis and  $24.81 \pm 2.32$  among eyes without



**Fig 2. 67 year old male with advanced POAG, with peripapillary retinoschisis.** (2A) Optic nerve photograph shows glaucomatous optic neuropathy. (2B) There is advanced visual field damage, with a mean deviation of -19.65 dB. (2C) OCT demonstrates peripapillary retinoschisis in the outer nuclear layer and outer plexiform layer (C; white arrows) on two horizontal raster scans through the optic nerve. (2D) Circumpapillary RNFL thickness map shows retinoschisis nasally.

<https://doi.org/10.1371/journal.pone.0182816.g002>



**Fig 3. 54 year old male with pigmentary OAG with peripapillary retinoschisis and peripapillary atrophy.** (3A and 3B) Two horizontal OCT raster scans through the optic nerve head show splitting in the inner nuclear layer (white star) and outer plexiform layer (white arrows). There is a focal area of vitreopapillary traction at the temporal margin of the optic nerve head (yellow arrows). (3C) Humphrey visual field shows mild glaucomatous damage with a MD of -1.99 dB. (3D) Circumpapillary RNFL thickness map and section shows RNFL segmentation sparing the retinoschisis in the inner nuclear layer (white star), outer nuclear and outer plexiform layers (white arrow).

<https://doi.org/10.1371/journal.pone.0182816.g003>

**Table 1. Clinical characteristics of the eleven eyes of seven patients with peripapillary retinal splitting (schisis).**

	Age/ Gender	Diagnosis	Eye	IOP	Layers Involved	Prior Surgery/ Laser	Glaucoma Meds	Other ocular pathology	BCVA	CCT (um)	SEQ (D)	VF MD (dB)	VF PSD	Axial length (mm)	CD Ratio	PPA
1	54M	Pigmentary OAG	OD	15	INL, ONL, OPL	None	Latanoprost	None	20/20	574	-8.75	-1.99	1.88	NA	0.7	Yes
2	78M	POAG	OD	11	NFL, IPL, OPL, ONL	CE	Bimatoprost, Dorzolamide	None	20/20	513	-0.75	0.09	2.02	22.71	0.75	No
3	78M	POAG	OS	11	INL, ONL	CE	Bimatoprost, dorzolamide	None	20/30	511	-1.25	-2.19	2.19	22.76	0.75	No
4	66M	OHT	OD	19	NFL, INL, ONL	None	None	None	20/25	531	-7.75	-0.08	2.98	NA	0.4	Yes
5	66M	OHT	OS	19	NFL, INL, ONL	None	None	None	20/25	543	-8.50	-0.86	1.88	NA	0.3	Yes
6	69F	OHT	OD	17	NFL, ONL	None	Latanoprost	None	20/40	573	-3.50	-1.02	2.26	27.74	0.8	No
7	69F	OHT	OS	14	NFL, ONL	CE	Latanoprost	None	20/25	574	-4.50	-2.83	1.98	27.90	0.8	No
8	67M	CNAG	OS	17	INL, ONL, OPL	LPI, peripheral iridoplasty	Brimonidine, Brinzolamide, Latanoprost	None	20/20	577	+4.75	-1.53	2.49	NA	0.1	No
9	67M	POAG	OD	14	INL, ONL,	Trab, Tube Shunt	Brimonidine	None	20/30	536	-2.75	-21.03	11.47	NA	0.8	Yes
10	67M	POAG	OS	11	NFL, INL, ONL	Trab	None	None	20/30	544	-3.25	-19.65	13.42	NA	0.7	Yes
11	80F	POAG	OS	10	NFL, INL, ONL	Trab	Timolol, Latanoprost	None	20/70	470	-3.25	-27.61	8.98	25.01	0.9	Yes

POAG = primary open angle glaucoma; OHT = ocular hypertension; CNAG = chronic narrow angle glaucoma; CE = cataract extraction; LPI = laser peripheral iridotomy; trab = trabeculectomy; IOP = intraocular pressure; BCVA = best corrected visual acuity; CCT = central corneal thickness; SEQ = spherical equivalent; VF MD = visual field mean deviation; VF PSD = visual field pattern standard deviation; CD Ratio = cup disc ratio; PPA = peripapillary atrophy; NFL = nerve fiber layer; OPL = outer plexiform layer; ONL = outer nuclear layer; INL = inner nuclear layer.

<https://doi.org/10.1371/journal.pone.0182816.t001>

peripapillary schisis ( $p = 0.195$ ). Spherical equivalent (Dioptres) was  $-3.6 \pm 3.9$  in eyes with peripapillary schisis compared to  $-1.4 \pm 3.4$  in those without ( $p = 0.016$ ). Among the 11 eyes with peripapillary retinoschisis, only 2 eyes were highly myopic (spherical equivalent  $\leq -8.0$  diopters or axial length  $> 26.0$  mm). The other eyes were mildly to moderately myopic and there was one hyperope. Review of the characteristics of peripapillary retinoschisis between the hyperopic and the myopic eyes did not reveal a significant difference.

## Discussion

Using high density OCT scans, centered on the optic nerve, we observed peripapillary retinal splitting in 1.1% of 990 consecutive glaucoma and glaucoma suspect eyes examined in our clinic. We found that eyes with peripapillary schisis were significantly more myopic and more likely to have vitreopapillary traction compared to eyes without peripapillary schisis. However, there was no significant difference in age, IOP, visual field mean deviation, pattern standard

**Table 2. Distribution of diagnosis in eyes with and without peripapillary retinal splitting (schisis).**

Eyes with Peripapillary Schisis	n = 11 eyes
Primary Open Angle Glaucoma	5 (45.4%)
Glaucoma Suspect	4 (36.3%)
Primary Angle Closure Glaucoma	1 (9.1%)
Pigmentary Glaucoma	1 (9.1%)
Eyes without Peripapillary Schisis	n = 979 eyes
Primary Open Angle Glaucoma	481 (49.1%)
Glaucoma Suspect	162 (16.5%)
Chronic Narrow Angle Glaucoma	45 (4.6%)
Pseudoexfoliation Glaucoma	52 (5.3%)
Ocular Hypertension	120 (12.3%)
Uveitic Glaucoma	33 (3.4%)
Pigmentary Open Angle Glaucoma	27 (2.8%)
Other Secondary Open Angle Glaucoma	24 (2.5%)
Pigment Dispersion Syndrome	2 (0.6%)

<https://doi.org/10.1371/journal.pone.0182816.t002>

deviation, prevalence of beta-zone peripapillary atrophy, vertical cup disk ratio or axial length between the two groups.

Peripapillary and macular schisis has been described in POAG[13], sometimes associated with serous detachment in eyes with a large optic nerve head cup[14], pseudoexfoliation glaucoma [15] and normal tension glaucoma. [8] The differential diagnosis of splitting in our patients includes myopic foveoschisis, tractional retinoschisis and traction on the disc. While our cross-sectional study cannot determine true causality, there have been various theories put forward to explain the pathogenesis of peripapillary schisis in glaucomatous eyes.

Some postulate that the development of micro-holes in the thinned area of the ONH or RNFL may allow liquid vitreous to enter the retina. [5, 7, 16] The presence of the macular schisis in normal-tension glaucoma also supports the micro-holes theory. [8] There has also been suggestion of a communicating bridge connecting the optic disc and the macular retinoschisis region, contraction of the membrane tissue on the optic disc creating a cleft, an optic nerve anomalous excavation functioning similar to a bulb syringe, and entry of vitreous through the lamina cribrosa, which may also be influenced by alterations in pulse or in the intracranial pressure due to changes in posture. [2] [17] [18] [19] [20] None of the eyes in our series had evidence of optic pits on OCT which have been associated with schisis. [2] [21] [22]

The second theory is that increased IOP and optic nerve cupping also may play a role in the development of retinoschisis. [5] [8] [23] It has also been suggested that small changes in axial length accompanying fluctuations in IOP may lead to vitreous traction, with or without microscopic breaks in the inner retina and play a role in schisis formation which has been reported in chronically elevated IOP. [6] [5, 19, 23, 24] None of the eyes with peripapillary schisis had IOP >21, while among eyes without schisis, 104 eyes had IOP >21 at the time of imaging or on their preceding clinic visit. Overall, there was no difference in the IOP among eyes with and without peripapillary schisis. Similar to our results, Hwang et al. [11] did not find any differences in IOP or visual field indices prior to or at the time of peripapillary retinoschisis formation.

Thirdly, microscopic fractures in the ILM could provide a conduit between the posterior cortical vitreous and the subretinal space [25] leading to macular neurosensory detachments. However, in our series, all cases of schisis extended back to the nerve margins suggesting an optic disk origin rather than an underlying ILM abnormality.



Lastly, vitreomacular or vitreopapillary traction secondary to incomplete vitreous separation is another plausible mechanism. [1, 26–30] [31] In our series, 7/11 (63.6%) of the eyes with peripapillary schisis had evidence of adherent vitreous with traction at the nerve in contrast to 13/979 (1.33%) of eyes without peripapillary schisis, while 404/979 (41.3%) eyes without peripapillary schisis had vitreopapillary adhesion without traction. In a series of 13 POAG eyes with microcystic INL changes similar to schisis, Hasegawa et al reported that the vitreous was attached in the area of microcystic INL spaces in 76.9% of eyes. A partial PVD has also been shown to increase RNFL thickness measurements, suggesting the impact of vitreous traction on the inner retinal layers. [32]

With extension into the macula, retinoschisis can impact vision and may warrant surgical intervention. [29] [7] Spontaneous resolution of macular disease as a result of complete vitreous separation has been reported in 10% of eyes with vitreomacular traction [33, 34] and it is possible that a similar course may be seen in glaucomatous eyes with peripapillary schisis leading to an underdetection of schisis.

Ouyang et al described 278 eyes, from 144 normal volunteers (81 females, 63 males) with a mean age of 37.6 years (range 18–74; SD = 15.5 years) that were imaged with the 3DOCT-1000 (Topcon Corp., Tokyo, Japan) using the raster scan protocol with  $512 \times 128$  scans, centered at the optic nerve head to report the prevalence of juxtapapillary pigment epithelium detachments. [35] These were asymptomatic volunteers that had not been diagnosed with any ocular pathology. We performed a post-hoc analysis of this dataset and upon review of images, found that none of the eyes had peripapillary schisis. While this does not cohort is underpowered to serve as a control group and was significantly younger, and predominantly hispanic, the absence of peripapillary schisis is in contrast to the 1.1% prevalence in our glaucoma and glaucoma suspect group.

The prevalence of peripapillary retinoschisis in glaucoma patients and glaucoma suspects has several implications. Firstly, coexisting retinoschisis in a patient with glaucoma may interfere with the reliability of RNFL thickness analysis, as retinoschisis may mask glaucomatous RNFL thinning due to inclusion of the retinoschisis cavities by the RNFL segmentation algorithm causing artifactual thickening and this should be considered when interpreting RNFL thickness measurements. In a recent study of 19 glaucomatous eyes with retinoschisis, the RNFL thickness was 81.6 microns, significantly greater than 69.7 microns in a group of age and visual field mean deviation matched glaucomatous eyes. [11] It has been reported that peripapillary retinoschisis is most commonly located in the superior quadrant (63.2%), followed by the inferior quadrant (15.8%), nasal quadrant (10.5%) and the superior-nasal area (5.3%). Clinicians should pay attention to the OCT segmentation algorithm to manually verify accurate segmentation and identify any anomalous pathology in the retinal layers. Retinoschisis has been reported in all the inner retinal layers including the RNFL, GCL and IPL. RNFL thickness has been shown to return to baseline levels following resolution of retinoschisis. [11] In our series, all eyes had splitting in the ONL, 72.3% in INL, 63.6% in the RNFL and 27.3% in the OPL.

Secondly, glaucoma patients and glaucoma suspects with vitreous traction should be followed for the development of retinoschisis as this may impact the visual field in the absence of true glaucomatous progression. High density cross sectional raster images through the optic nerve can accurately identify vitreopapillary traction which may be missed on a 3.46 mm diameter circle RNFL scan.

Thirdly, this is also applicable to eyes with peripapillary atrophy as 54.5% of the eyes with schisis in our study had beta-zone peripapillary atrophy. Although we did not use enhanced depth imaging-OCT, it is also possible that the peripapillary atrophy observed was along the



spectrum of peripapillary choroidal cavitation which has been associated with schisis. [36] [37, 38] [39] [40, 41]

The majority of patients in our series with peripapillary retinoschisis had POAG (42.9%) followed by glaucoma suspects (28.6%), chronic narrow angle glaucoma (14.3%) and pigmentary glaucoma (14.3%), suggesting that schisis may occur across the spectrum of the glaucoma continuum and not just in those eyes with advanced disease. Unlike a typical area of retinoschisis found in the peripheral retina, the schisis associated with optic disc pits does not cause an absolute scotoma [19] and thus may not cause changes in the visual field, making detection using imaging more important.

Limitations of our study include its retrospective nature, lack of an adequately powered non-disease group that precludes us from calculating the prevalence and predictive value of peripapillary schisis in glaucoma and glaucoma suspects, and inability to determine true causality in this cross sectional study.

Unique to our investigation, to the best of our knowledge, is the review of a large series of consecutive glaucoma patients and glaucoma suspects who underwent high resolution cross sectional OCT raster scans to detect peripapillary schisis. Based on these data, we report peripapillary schisis in 1.1% of a large cohort of glaucoma patients and glaucoma-suspects. Evidence of adherent vitreous with traction and peripapillary atrophy was found in a majority of the eyes with schisis. Further long term prospective studies analyzing the clinical course and characteristics of peripapillary retinoschisis are warranted to better elucidate the clinical significance of this pathology.

## Supporting information

**S1 Table. Supporting data file contains the information of the 990 glaucoma and glaucoma suspect eyes including the 11 eyes with schisis.**  
(XLSX)

## Acknowledgments

Meeting Presentation: This manuscript was presented in part as a poster at the Association for Research in Vision and Ophthalmology Annual Meeting in May 2014 and the American Ophthalmological Society Annual Meeting in May 2014.

## Author Contributions

**Conceptualization:** Dilraj S. Grewal, Angelo P. Tanna.

**Data curation:** Daniel J. Merlau, Pushpanjali Giri, Angelo P. Tanna.

**Investigation:** Amani A. Fawzi, Angelo P. Tanna.

**Methodology:** Angelo P. Tanna.

**Project administration:** Angelo P. Tanna.

**Supervision:** Lee M. Jampol, Angelo P. Tanna.

**Writing – original draft:** Dilraj S. Grewal.

**Writing – review & editing:** Dilraj S. Grewal, Marion R. Munk, Amani A. Fawzi, Lee M. Jampol, Angelo P. Tanna.

## References

1. Postel EA, Pulido JS, McNamara JA, Johnson MW. The etiology and treatment of macular detachment associated with optic nerve pits and related anomalies. *Transactions of the American Ophthalmological Society*. 1998; 96:73–88; discussion -93. Epub 1999/06/09. PMID: [10360283](#).
2. Johnson TM, Johnson MW. Pathogenic implications of subretinal gas migration through pits and atypical colobomas of the optic nerve. *Archives of ophthalmology*. 2004; 122(12):1793–800. Epub 2004/12/15. <https://doi.org/10.1001/archophth.122.12.1793> PMID: [15596582](#).
3. Georgalas I, Ladas I, Georgopoulos G, Petrou P. Optic disc pit: a review. *Graefe's archive for clinical and experimental ophthalmology = Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie*. 2011; 249(8):1113–22. <https://doi.org/10.1007/s00417-011-1698-5> PMID: [21638030](#).
4. Chang S, Gregory-Roberts E, Chen R. Retinal detachment associated with optic disc colobomas and morning glory syndrome. *Eye*. 2012; 26(4):494–500. Epub 2012/01/14. <https://doi.org/10.1038/eye.2011.354> PMID: [22241012](#).
5. Hollander DA, Barricks ME, Duncan JL, Irvine AR. Macular schisis detachment associated with angle-closure glaucoma. *Archives of ophthalmology*. 2005; 123(2):270–2. <https://doi.org/10.1001/archophth.123.2.270> PMID: [15710832](#).
6. Kahook MY, Noecker RJ, Ishikawa H, Wollstein G, Kagemann L, Wojtkowski M, et al. Peripapillary schisis in glaucoma patients with narrow angles and increased intraocular pressure. *American journal of ophthalmology*. 2007; 143(4):697–9. <https://doi.org/10.1016/j.ajo.2006.10.054> PMID: [17386284](#).
7. Zumbro DS, Jampol LM, Folk JC, Olivier MM, Anderson-Nelson S. Macular schisis and detachment associated with presumed acquired enlarged optic nerve head cups. *Am J Ophthalmol*. 2007; 144(1):70–4. <https://doi.org/10.1016/j.ajo.2007.03.027> PMID: [17493573](#).
8. Zhao M, Li X. Macular retinoschisis associated with normal tension glaucoma. *Graefe's archive for clinical and experimental ophthalmology = Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie*. 2011; 249(8):1255–8. <https://doi.org/10.1007/s00417-011-1668-y> PMID: [21455774](#).
9. Hasegawa T, Akagi T, Yoshikawa M, Suda K, Yamada H, Kimura Y, et al. Microcystic Inner Nuclear Layer Changes and Retinal Nerve Fiber Layer Defects in Eyes with Glaucoma. *PloS one*. 2015; 10(6):e0130175. <https://doi.org/10.1371/journal.pone.0130175> PMID: [26066021](#).
10. Lee EJ, Kim TW, Kim M, Choi YJ. Peripapillary retinoschisis in glaucomatous eyes. *PLoS One*. 2014; 9(2):e90129. <https://doi.org/10.1371/journal.pone.0090129> PMID: [24587238](#).
11. Hwang YH, Kim YY, Kim HK, Sohn YH. Effect of peripapillary retinoschisis on retinal nerve fibre layer thickness measurement in glaucomatous eyes. *The British journal of ophthalmology*. 2014. Epub 2014/02/08. <https://doi.org/10.1136/bjophthalmol-2013-303931> PMID: [24505035](#).
12. Silverman AL, Hammel N, Khachatryan N, Sharpsten L, Medeiros FA, Girkin CA, et al. Diagnostic Accuracy of the Spectralis and Cirrus Reference Databases in Differentiating between Healthy and Early Glaucoma Eyes. *Ophthalmology*. 2015. <https://doi.org/10.1016/j.ophtha.2015.09.047> PMID: [26526632](#).
13. Hubschman JP, Reddy S, Kaines A, Law S. Nasal Retinoschisis Associated with Glaucoma. *Ophthalmic surgery, lasers & imaging: the official journal of the International Society for Imaging in the Eye*. 2010; 1–4. <https://doi.org/10.3928/15428877-20100215-60> PMID: [20337311](#).
14. Farjad H, Besada E, Frauens BJ. Peripapillary Schisis with Serous Detachment in Advanced Glaucoma. *Optometry and vision science: official publication of the American Academy of Optometry*. 2010. <https://doi.org/10.1097/OPX.0b013e3181d1dad5> PMID: [20125057](#).
15. Ornek N, Buyuktortop N, Ornek K. Peripapillary and macular retinoschisis in a patient with pseudoexfoliation glaucoma. *BMJ case reports*. 2013; 2013. <https://doi.org/10.1136/bcr-2013-009469> PMID: [23729710](#).
16. Xin D, Talamini CL, Raza AS, de Moraes CG, Greenstein VC, Liebmann JM, et al. Hypodense regions (holes) in the retinal nerve fiber layer in frequency-domain OCT scans of glaucoma patients and suspects. *Investigative ophthalmology & visual science*. 2011; 52(10):7180–6. Epub 2011/07/28. <https://doi.org/10.1167/iovs.11-7716> PMID: [21791587](#).
17. Rutledge BK, Puliafito CA, Duker JS, Hee MR, Cox MS. Optical coherence tomography of macular lesions associated with optic nerve head pits. *Ophthalmology*. 1996; 103(7):1047–53. Epub 1996/07/01. PMID: [8684793](#).
18. Mavrikakis E, Lam WC. Macular schisis and detachment secondary to large optic nerve head cup: a newly recognized syndrome amenable to vitrectomy. *Acta ophthalmologica*. 2011; 89(1):95–6. Epub 2009/06/25. <https://doi.org/10.1111/j.1755-3768.2009.01553.x> PMID: [19549105](#).
19. Spaide RF, Costa DL, Huang SJ. Macular schisis in a patient without an optic disk pit optical coherence tomographic findings. *Retina*. 2003; 23(2):238–40. Epub 2003/04/23. PMID: [12707606](#).

20. Takashina S, Saito W, Noda K, Katai M, Ishida S. Membrane tissue on the optic disc may cause macular schisis associated with a glaucomatous optic disc without optic disc pits. *Clinical ophthalmology*. 2013; 7:883–7. <https://doi.org/10.2147/OPTH.S42085> PMID: 23690678.
21. Coll GE, Chang S, Flynn TE, Brown GC. Communication between the subretinal space and the vitreous cavity in the morning glory syndrome. *Graefe's archive for clinical and experimental ophthalmology = Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie*. 1995; 233(7):441–3. Epub 1995/07/01. PMID: 7557510.
22. Friberg TR, McLellan TG. Vitreous pulsations, relative hypotony, and retrobulbar cyst associated with a congenital optic pit. *American journal of ophthalmology*. 1992; 114(6):767–9. Epub 1992/12/15. PMID: 1463050.
23. Lampert PW, Vogel MH, Zimmerman LE. Pathology of the optic nerve in experimental acute glaucoma. Electron microscopic studies. *Investigative ophthalmology*. 1968; 7(2):199–213. Epub 1968/04/01. PMID: 4966696.
24. Pilon A, Newman T, Messner LV. Neurosensory detachment arising from a fractured inner-limiting membrane secondary to chronically elevated intraocular pressure. *Optom Vis Sci*. 2006; 83(7):415–20. <https://doi.org/10.1097/01.opx.0000218430.40132.68> PMID: 16840866.
25. Johnson MW. Perifoveal vitreous detachment and its macular complications. *Transactions of the American Ophthalmological Society*. 2005; 103:537–67. PMID: 17057817.
26. Hwang YH, Kim YY. Peripapillary Retinal Nerve Fiber Layer Thickening Associated with Vitreopapillary Traction. *Seminars in ophthalmology*. 2013. Epub 2013/11/01. <https://doi.org/10.3109/08820538.2013.833257> PMID: 24171792.
27. Hirakata A, Hida T, Wakabayashi T, Fukuda M. Unusual posterior hyaloid strand in a young child with optic disc pit maculopathy: intraoperative and histopathological findings. *Japanese journal of ophthalmology*. 2005; 49(3):264–6. <https://doi.org/10.1007/s10384-004-0185-5> PMID: 15944837.
28. Besada E, Barr R, Schatz S, Brewer C. Vitreal pathogenic role in optic pit foveolar retinoschisis and central serous retinopathy. *Clinical & experimental optometry: journal of the Australian Optometrical Association*. 2003; 86(6):390–8. PMID: 14632616.
29. Hirakata A, Inoue M, Hiraoka T, McCuen BW 2nd. Vitrectomy without laser treatment or gas tamponade for macular detachment associated with an optic disc pit. *Ophthalmology*. 2012; 119(4):810–8. <https://doi.org/10.1016/j.ophtha.2011.09.026> PMID: 22218142.
30. Haruta M, Kamada R, Umeno Y, Yamakawa R. Vitrectomy for optic disc pit-associated maculopathy with or without preoperative posterior vitreous detachment. *Clinical ophthalmology*. 2012; 6:1361–4. <https://doi.org/10.2147/OPTH.S34962> PMID: 22973089.
31. Gandorfer A, Rohleder M, Grosselfinger S, Haritoglou C, Ulbig M, Kampik A. Epiretinal pathology of diffuse diabetic macular edema associated with vitreomacular traction. *American journal of ophthalmology*. 2005; 139(4):638–52. Epub 2005/04/06. <https://doi.org/10.1016/j.ajo.2004.11.035> PMID: 15808159.
32. Batta P, Engel HM, Shrivastava A, Freeman K, Mian U. Effect of partial posterior vitreous detachment on retinal nerve fiber layer thickness as measured by optical coherence tomography. *Archives of ophthalmology*. 2010; 128(6):692–7. Epub 2010/06/16. <https://doi.org/10.1001/archophthalmol.2010.99> PMID: 20547945.
33. Hikichi T, Yoshida A, Trempe CL. Course of vitreomacular traction syndrome. *American journal of ophthalmology*. 1995; 119(1):55–61. Epub 1995/01/01. PMID: 7825690.
34. Kusaka S, Saito Y, Okada AA, Sasamoto M, Hayashi A, Ohji M, et al. Optical coherence tomography in spontaneously resolving vitreomacular traction syndrome. *Ophthalmologica Journal international d'ophtalmologie International journal of ophthalmology Zeitschrift fur Augenheilkunde*. 2001; 215(2):139–41. Epub 2001/03/13. PMID: 11244348.
35. Ouyang Y, Heussen FM, Keane PA, Pappuru RK, Sadda SR, Walsh AC. Juxtapapillary pigment epithelium detachment observed in asymptomatic participants using optical coherence tomography. *Investigative ophthalmology & visual science*. 2013; 54(2):1144–9. <https://doi.org/10.1167/iovs.12-9903> PMID: 23322565.
36. Akimoto M, Akagi T, Okazaki K, Chihara E. Recurrent macular detachment and retinoschisis associated with intrachoroidal cavitation in a normal eye. *Case reports in ophthalmology*. 2012; 3(2):169–74. Epub 2012/06/09. <https://doi.org/10.1159/000339292> PMID: 22679435.
37. Shimada N, Ohno-Matsui K, Nishimuta A, Tokoro T, Mochizuki M. Peripapillary changes detected by optical coherence tomography in eyes with high myopia. *Ophthalmology*. 2007; 114(11):2070–6. Epub 2007/06/05. <https://doi.org/10.1016/j.ophtha.2007.01.016> PMID: 17543388.
38. Shimada N, Ohno-Matsui K, Iwanaga Y, Tokoro T, Mochizuki M. Macular retinal detachment associated with peripapillary detachment in pathologic myopia. *International ophthalmology*. 2009; 29(2):99–102. Epub 2007/11/24. <https://doi.org/10.1007/s10792-007-9174-2> PMID: 18034213.

39. Park SC, De Moraes CG, Teng CC, Tello C, Liebmann JM, Ritch R. Enhanced depth imaging optical coherence tomography of deep optic nerve complex structures in glaucoma. *Ophthalmology*. 2012; 119(1):3–9. <https://doi.org/10.1016/j.ophtha.2011.07.012> PMID: 21978593.
40. Fellman RL, Grover DS. Myopic peripapillary sinkhole: prolapse of retinal nerve fiber layer and posterior vitreous into a sclerochoroidal hollow causing peripapillary choroidal thickening and cavitation. *Archives of ophthalmology*. 2012; 130(9):1220–1. Epub 2012/09/12. <https://doi.org/10.1001/archophthalmol.2012.441> PMID: 22965607.
41. Freund KB, Mulkamala SK, Cooney MJ. Peripapillary choroidal thickening and cavitation. *Archives of ophthalmology*. 2011; 129(8):1096–7. Epub 2011/08/10. <https://doi.org/10.1001/archophthalmol.2011.208> PMID: 21825199.